

levels varied inversely, with the highest occurring during the early morning hours. Histamine levels did not rise comparably in subjects without asthma. Results of these studies imply that some breakdown in the homeostatic mechanism for controlling bronchomotor tone occurs in persons with asthma. Because antihistamines are not regularly effective against nocturnal asthma, other mediators of bronchoconstriction need to be determined before this aspect of asthma can be treated.

HAROLD S. NOVEY, MD

REFERENCES

- Connolly CK: Diurnal rhythms in airway obstruction. *Br J Dis Chest* 73:357-366, Oct 1979
 Hetzel MR, Clark TJ: Does sleep cause nocturnal asthma? *Thorax* 34:749-754, Dec 1979
 Barnes P, FitzGerald G, Brown M, et al: Nocturnal asthma and changes in circulating epinephrine, histamine, and cortisol. *N Engl J Med* 303:263-267, Jul 31, 1980

Physical Urticarias: An Important Cause of Hives

THE PHYSICAL URTICARIAS are a group of diseases in which certain stimuli applied to the skin cause hives and, occasionally, respiratory symptoms. At least seven distinct forms of dermal stimuli can cause urticaria: heat, cold, sunlight, trauma, prolonged pressure, vibration and water. Heat, cold and solar urticaria, however, each exist in more than one form. Physical urticarias are usually acquired conditions, although there are two rare inherited forms—familial cold urticaria and vibratory urticaria. Generalized heat urticaria (cholinergic urticaria) and trauma urticaria (dermatographism) are relatively common, the latter occurring in 1 percent to 5 percent of the population. The mechanisms of most physical urticarias are poorly understood, although mast cell mediators are clearly involved in several of the forms. There is no association with other atopic disease, but immunoglobulin E antibodies may play a role in dermatographism and in some forms of acquired cold urticaria and solar urticaria.

The diagnosis of physical urticaria can usually be made by taking a careful history and carrying out appropriate challenge tests. Laboratory tests are of use only in excluding other conditions. Localized heat urticaria and primary acquired cold urticaria can be determined by exposure of an extremity to hot water or an ice cube, respectively. Dermatographism can be determined by stroking the chest or back with a blunt object, and aquagenic urticaria can be detected by application of distilled water to the skin. Solar urticaria can

be induced by exposure to sunlight or ultraviolet light of specified wavelengths and vibratory urticaria by skin massage or vibration. Familial cold urticaria differs from acquired cold urticaria by a delayed response to cold (negative ice cube test), positive family history and systemic findings (leukocytosis, fever and arthralgias) at the time of the urticaria. Pressure urticaria usually occurs after prolonged pressure, and the wheals characteristically develop in association with tight clothing, especially at the waistline, shoulders, ankles or feet.

Cholinergic urticaria has a distinctive appearance of tiny wheals on a large erythematous base over the trunk, and it may become generalized with resulting bronchospasm. It is thought to be due to a dermal hypersensitivity to released acetylcholine and, thus, is provoked by exercise or factors which induce sweating (such as stress, emotion or heat). The intradermal test with methacholine has not been shown to be reliable for diagnosis of this disease.

Hypovolemic shock can sometimes occur with physical urticarias if the stimulus is intense or occurs over a large surface area. This is seen most often in acquired cold urticaria when there is a sudden, diffuse exposure to cold such as when one plunges into cold water. Patients with recurrent urticaria or unexplained anaphylaxis should be questioned about physical urticarias, and provocative testing should be done when appropriate.

Treatment of physical urticarias consists of avoidance of the causative physical stimulus, and administration of antihistamine drugs. Solar urticaria may benefit from appropriate-spectrum sunscreens. Steroids are rarely required.

JOHN R. TENCATI, MD

REFERENCES

- Kaplan AP: Mediators of urticaria and angioedema. *J Allergy Clin Immunol* 60:324-332, Nov 1977
 Soter NA, Wasserman SI, Austen KF, et al: Release of mast-cell mediators and alterations in lung function in patients with cholinergic urticaria. *N Engl J Med* 302:604-608, Mar 13, 1980
 Soter NA, Wasserman SI: Physical urticaria/angioedema: An experimental model of mast cell activation in humans. *J Allergy Clin Immunol* 66:358-365, Nov 1980

Respiratory Symptoms in Children Whose Parents Smoke

EPIDEMIOLOGICAL STUDIES have shown that upper and lower respiratory symptoms are more likely to occur in children whose parents smoke. Survey findings from the United States, Great Britain and Israel indicate that such symptoms occur most commonly if both parents smoke, less commonly

if one parent smokes, and least often if there is no smoking in the household. It was found that in families in which one parent smoked, only smoking by the mother produced a significant effect on the child. The child's symptoms were directly correlated with the number of cigarettes smoked by the mother as well as with maternal respiratory symptoms, but not with birth weight, birth order, social class or family size.

The symptom most commonly observed in infants less than 1 year of age, whose parents smoke, is cough, particularly during daytime hours, whereas older children generally have persistent wheezing in addition to coughing. Dyspnea and sputum production as well as upper respiratory symptoms that suggest recurrent colds or allergic rhinitis occur at all ages. Pulmonary function is inversely correlated with cigarette consumption by parents. There is no increased incidence of croup or bronchiolitis.

Although a familial genetic susceptibility to respiratory illness cannot be ruled out by such studies, it is more likely that the well-recognized

irritant properties of cigarette smoke ("passive smoking") are the major factor in producing these ailments. Findings of several studies show a positive correlation of respiratory symptoms in infants and children with respiratory symptoms in their parents who smoke, which suggests cross-infection within the household as a possible mechanism. However, it is more likely that symptoms of both the children and adults are a reflection of the amount of exposure to smoke.

In the diagnosis of chronic coughing in infants and persistent wheezing in children, parental smoking must be considered to avoid possible misdiagnosis as recurrent colds, hay fever, asthma, bronchitis or pneumonia.

ABBA I. TERR, MD

REFERENCES

- Colley JR, Holland WW, Corkhill RT: Influence of passive smoking and parental phlegm on pneumonia and bronchitis in early childhood. *Lancet* 2:1031-1034, Nov 2, 1974
- Harlap S, Davies AM: Infant admissions to hospital and maternal smoking. *Lancet* 1:529-532, Mar 30, 1974
- Lebowitz MD, Burrows B: Respiratory symptoms related to smoking habits of family adults. *Chest* 69:48-50, Jan 1976
- Weiss ST, Tager IB, Speizer FE, et al: Persistent wheeze: Its relation to respiratory illness, cigarette smoking, and level of pulmonary function in a population sample of children. *Am Rev Respir Dis* 122:697-707, Nov 1980

ADVISORY PANEL TO THE SECTION ON ALLERGY

ABBA I. TERR, MD

Advisory Panel Chairman

CMA Scientific Board Representative

Section Editor

San Francisco

NED J. WHITCOMB, MD
CMA Section Chairman
Carmichael

ROGER M. KATZ, MD
CMA Section Secretary
Los Angeles

NORMAN LEWISTON, MD
Stanford University

STEPHEN M. NAGY, JR, MD
University of California, Davis

HAROLD S. NOVEY, MD
University of California, Irvine

KHING ONG, MD
University of California, Irvine

GILDON BEALL, MD
University of California, Los Angeles

DOUGLAS HEINER, MD
University of California, Los Angeles

ROBERT HAMBURGER, MD
University of California, San Diego

OSCAR L. FRICK, MD
Section Editor
University of California, San Francisco

ZACK HADDAD, MD
University of Southern California
Los Angeles

JAMES P. KEMP, MD
San Diego

ELI MELTZER, MD
San Diego

NORMAN D. McCANN, MD
Chico

MITCHELL SOLLOD, MD
San Francisco

ALAN BECKER, MD
Tarzana

ALAN SINGER, MD
Torrance

THEODORE CHU, MD
San Jose

DANIEL GORENBERG, MD
San Bernardino